

L1 ANSWER 1 OF 1 WPIX COPYRIGHT 2008 THE THOMSON CORP on STN  
 ACCESSION NUMBER: 1987-314145 [45] WPIX  
 DOC. NO. CPI: C1987-133624 [21]  
 TITLE: N-substd. 3-amino:thieno-(2,3-B)-pyridine- 2-carboxamide  
 cpds. production - e.g. reaction of corresp. 2-carboxylic  
 acid ester(s) with amine(s)  
 DERWENT CLASS: B02; E13  
 INVENTOR: BOHM N; LABAN G; LEISTNER S; LOHMANN D; VIEWEG H; WAGNER  
 G  
 PATENT ASSIGNEE: (UYLE-C) UNIV KARL MARX LEIPZIG  
 COUNTRY COUNT: 1  
 PATENT INFO ABBR.:

PATENT NO	KIND DATE	WEEK	LA	PG	MAIN IPC
DD 247002	A	19870624 (198745)*	DE	5[0]	<--

APPLICATION DETAILS:

PATENT NO	KIND	APPLICATION	DATE
DD 247002 A		DD 1986-287347	19860226

PRIORITY APPLN. INFO: DD 1986-287347 19860226

AN 1987-314145 [45] WPIX

AB DD 247002 A UPAB: 20050426

Production of N-substd. 3-amino-thieno (2,3-b)pyridine-2-carboxamide derivs. of formula (I) (many of which are new), comprises (A) reacting corresp. 2-carboxylic acid ester (II) with a primary or secondary amine. (IV) at boiling temperature, or (B) reacting a 2-(3-cyano-2-pyridylthio) acetic acid ester (IV) with a functionalised aliphatic amino, or (C) reacting a 2-mercapto-pyridine-3-carbonitrile (V) with a 2-halo-acetamide (VI) in alkanolic solution in the presence of alkali alcoholate.

USE - The products are intermediates for biologically active heterocyclic cpds. - In an example, 4,6-Dimethyl-2-mercaptopyridine-3-carbonitrile (16.4g) in IM ethanolic KOH (350 ml) is reacted with chloroacetic acid (10g) in water (20ml) to give 3-amino-4,6-dimethyl- thieno(2,3-b)pyridine-2-carboxylic acid potassium salt (m. pt. 360 deg.) in 85% yield. This is dissolved in water and acidified with acetic acid. The resulting free acid is dried in a vacuum desiccator are KOH/H2SO4. The resulting free acid (2.2g) and p-nitrophenol(1.8g) in absolute pyridine (35ml) are treated with dicyclohexylcarbodiimide (2.2g), reacted 20 hrs. at room temperature, and worked up to give the p-nitrophenyl ester (m.pt. 252-6 deg.) in 60% yield. Nitrophenyl ester (1-3g) and 2-diethylamino-ethylamine (1.5g) are refluxed 2 hrs. in absolute dioxane (15ml), then worked up to give N-(2-diethylaminoethyl) -3-amino-4,6-dimethyl-thieno (2,3-b)pyridine-2-carboxamide, m.pt. 130 deg. (from aqueous EtOH) in 72% yield.